

REVIEW

Tension pneumothorax—time for a re-think?

S Leigh-Smith, T Harris

Emerg Med J 2005;22:8–16. doi: 10.1136/emj.2003.010421

This review examines the present understanding of tension pneumothorax and produces recommendations for improving the diagnostic and treatment decision process.

Tension pneumothorax (TPT) is an uncommon disease with a malignant course leading to death if untreated.^{1,2} It is most commonly encountered in prehospital trauma care, emergency departments, and intensive care units (ICUs).³ Resuscitation and trauma courses usually illustrate a patient in extremis and assume that the clinical diagnosis is straightforward and the response to needle chest decompression is rapid and reliable.^{4,5} However, this might not be the case in real life. Texts differ when describing the diagnostic symptoms and signs^{6–9} and there are several case reports of diagnostic difficulty or missed diagnosis because of an absence of “classic” signs.^{1,10,11} Lack of chest signs along with poor correlation between the signs present and those picked up by experienced physicians have been specifically noted.^{12–14} There have also been multiple reports of ineffective needle decompression with adequate treatment only being achieved with tube thoracostomy.^{15–22}

This review examines the present understanding of tension pneumothorax and produces recommendations for improving the diagnostic and treatment decision process.

METHODOLOGY

An electronic search was performed without use of filters, on Medline (1966–2003), (once with Pubmed and once with Ovid interface). The terms “pneumothorax/*” (PNEUMOTHORAX) and “tension\$/*” were searched separately, then combined to identify relevant papers, which once obtained were hand searched, with further references being obtained from these. Original papers were systematically reviewed for pathophysiology, predefined clinical features, and treatment options. Both authors performed this process independently, writing separate reviews, which were then collated to achieve a consensus article.

TENSION PNEUMOTHORAX—DEFINITION

TPT may be said to occur when a one way valve communicates with the pleural space,^{5,23} but this in fact describes an injury with the potential to tension. We found a variety of definitions of tension pneumothorax (box 1).

Definitions using intrapleural pressure (IPP) are more accurate than clinical definitions and

the definition of “positive IPP throughout the respiratory cycle” was used in early animal experiments. However, in the awake subject, the IPP must be less than atmospheric pressure during part of the respiratory cycle if the pneumothorax is to continue to develop, as this will only occur if air continues to enter the pleural cavity. This led to the definition of an “expiratory tension pneumothorax.”^{33,34} It is therefore important to consider that in any TPT (in an awake subject) there is a range from an IPP that is only just positive at end expiration to an IPP that is positive throughout the respiratory cycle.³⁵

In some definitions supra-physiological IPP occurs before the defined TPT exists. This may have limited clinical relevance, however it makes the literature difficult to interpret as papers may be referring to different stages in development of the condition.

INCIDENCE

The incidence of TPT varies with the population studied and is not well established—often reflecting disease suspicion rather than true incidence. TPT was confirmed (hiss of air on decompression) in 5.4% of major trauma patients, (64% of whom were ventilated) treated by prehospital care doctors in London.³⁶ Other studies have reported needle decompression rates (as a surrogate for potential TPT) in the prehospital environment varying from 0.7% to 30%.^{32,37,38}

Earlier postmortem studies on patients dying in ICUs show rates of undiagnosed TPT ranging from 1.1% to 3.8%.^{25,26} Missed diagnosis was more probable if ventilation or cardiopulmonary resuscitation had occurred.

In ventilated patients TPT is more likely if simple pneumothorax diagnosis is delayed.^{39,27} TPT would also seem to be more serious in ventilated patients reaching 91% mortality rates in one series.²⁹

PATHOPHYSIOLOGY

In the awake patient IPP is sub-atmospheric (during inspiration and expiration) resulting from the elastic recoil forces of the lung and in a healthy awake subject it ranges from –5 to –8 cm H₂O. A forced maximal inspiratory effort can achieve an IPP of –80 cm H₂O and this is important in the awake patient trying to overcome the effects of increasing IPP as a TPT develops.

Abbreviations: TPT, tension pneumothorax; ICU, intensive care unit; IPP, intrapleural pressure; CVP, central venous pressure

See end of article for authors' affiliations

Correspondence to:
Mr S Leigh-Smith, c/o
Institute of Naval
Medicine, Monckton
House, Alverstoke,
Gosport, Hampshire PO12
2DL, UK;
simonlsuk@aol.com

Accepted for publication
6 January 2004

Box 1 Different definitions of tension physiology in a patient with a pneumothorax

Presence of:

- severe clinical manifestations²⁴
- hiss of air on thoracic needle decompression^{24 25}
- mediastinal shift
 - on chest radiograph at postmortem examination²⁶
 - +/- diaphragmatic depression on chest radiography²⁴
 - diagnosed clinically or on chest radiography in a ventilated patient²⁷
- haemodynamic compromise with
 - haemodynamic improvement and release of gas on tube thoracostomy²⁸
 - opposite mediastinal shift on chest radiography²⁸
 - expanding pneumothorax³⁰
- ipsilateral intrapleural pressure
 - higher than atmospheric pressure^{31 32}
 - positive throughout the respiratory cycle³⁰
 - positive during expiration^{33 34}

In contrast with this the paralysed, sedated, and ventilated patient has gas delivered under supra-physiological positive pressure. Inspiratory pressures are often in excess of +20 cm H₂O along with either physiological expiratory alveolar pressures or positive end expiratory pressure levels upwards of +5 cm H₂O.

Between the two extremes of awake and ventilated patients lie various levels of sedation and ventilatory support modes commonly used in ICUs. However, for simplicity we have assigned the evidence to two groups—awake (spontaneously ventilating) or ventilated.

Pneumothorax develops secondary to a breach in the visceral, parietal, or mediastinal pleura and is termed spontaneous (simple, idiopathic), traumatic, or iatrogenic depending on the precipitant. It is termed primary when in the absence of underlying clinically apparent respiratory disease and secondary if resultant from underlying pulmonary pathology.

If the pleural defect functions as a one way valve, air enters the pleural cavity on inspiration but is unable to exit on expiration, leading to increasing ipsilateral IPP. This will cause further lung collapse, chest wall expansion, diaphragmatic depression, and (dependent on mediastinal distensibility) contralateral lung compression.

ANIMAL MODELS—AWAKE

Animal experiments use intrapleural catheters to induce expanding pneumothoraces and early experiments in awake dogs noted a significant fall in cardiac output,^{40 41} which was most pronounced in the anaesthetised animals.⁴² This led to the theory of thoracic vein compression, right heart compression, and great vessel kinking causing reduced cardiac filling.^{40 41} It was however noted that—unlike humans—the dog's mediastinum was flimsy and unable to maintain a pressure gradient between the two pleural spaces,^{42 43} along with being perforate, resulting in unilaterally introduced air spreading bilaterally.^{40 44} These differences permit more direct transmission of IPP to central structures.³ Despite this difference the theory of great vessel kinking as a cause of cardiovascular collapse persists in studies and textbooks.

Later studies used goats, swine, and sheep, which—like humans—have an intact and fixed mediastinum.^{30 33 45 46} In one of these studies the animals were lightly anaesthetised (but spontaneously ventilating),³⁰ and in all others they were fully awake.^{33 45 47} These animals were able to maintain negative contralateral IPP, even after significant positive ipsilateral IPP occurred in the end expiratory phase^{33 47} and sometimes throughout the whole respiratory cycle.³⁰

One study induced bilateral TPT in dogs⁴⁵ to overcome the dog model validity problems mentioned above, but this may negate any contralateral thoracic compensatory mechanisms, which occur with unilateral TPT.

These studies showed a variety of progressive compensatory mechanisms (box 2) before final collapse with some animals tolerating extreme degrees of “tension” very well.^{30 45}

Minute volume and ventilation (to normocapnia) were preserved until a very late stage despite universal findings of progressive and severe hypoxia,^{30 33 45 47} which suggests pulmonary shunting and parenchymal collapse as the main causes of hypoxia.^{30 45 47}

Despite falling stroke volumes cardiac output was almost universally preserved throughout^{30 33 45 47} with one study noting a 19% increase in BP.⁴⁷ In all these studies just two animals (both immature monkeys with mobile mediastinums) developed hypotension immediately before death from respiratory arrest, but cardiac output was still 70% normal at this stage.³⁰ Preservation of cardiac output is through a progressive tachycardia, incomplete transmission of positive IPP to the mediastinum, and increasingly negative contralateral intrathoracic pressure preserving the venous return.^{30 33 47}

There was minimal evidence of right sided venous obstruction with only bilateral TPTs causing a linear rise in central venous pressure (CVP).⁴⁵ Only one of the three unilateral TPT studies found increased superior and inferior vena caval pressures and these occurred in the agonal stages of respiration.³⁰ Despite this there was no evidence of one specific point of obstruction or kinking in the circulation³⁰ and it was felt that unilateral tension does not mechanically compromise cardiac output.^{30 33}

The dominant physiological feature during decompensation was progressive respiratory failure with death from respiratory not cardiovascular arrest.^{30 33} This was postulated to be attributable to; myocardial hypoxia, central hypoxemia with hypoxic suppression of the respiratory centre, restrictive ventilatory defect, decreased efficiency of hypoxic intercostal muscles at the hyperinflated chest wall, and depression of the ipsilateral hemi-diaphragm from positive IPP.^{30 33 47}

Box 2 Compensatory mechanisms during tension pneumothorax in awake animals^{30 33 45 47}

Respiratory

- Increasing respiratory rate
- Increasing tidal volume
- Contralateral
 - increasingly negative IPP excursions
 - increasing chest wall expansion

Cardiac

- Tachycardia

ANIMAL MODELS—VENTILATED

The effect of raised IPP is very different if the animal is ventilated and sedated, as the cardiorespiratory compensatory mechanisms (box 2) are impaired. Also, during ventilation increased alveolar pressure may obstruct cardiac output, and if alveolar pressure rises above pulmonary venous (or pulmonary arterial) pressure cardiac output will be reduced.⁴⁸

Studies of increasing IPP in ventilated sheep and swine support this suggestion, along with showing progressive hypoxia as the earliest sign, and an immediate rise in CVP.^{46 49} Hypotension and tachycardia occurred after pneumothorax of 47% total lung capacity and were progressive, being well tolerated up until eventual and sudden cardiovascular collapse at pneumothorax of roughly 90% total lung capacity. Death occurred soon after at which point SpO₂ was <50%, and CVP equalled IPP suggesting complete occlusion of venous return at this pre-terminal stage.⁴⁹ Other findings were; V/Q mismatch, alveolar compression, and hypercapnia.^{46 49} These results suggest that a more rapid deterioration and early reduction in cardiac output is expected in ventilated patients.^{3 50}

TENSION PNEUMOTHORAX IN HUMANS

Awake patients

Box 3 summarises the symptoms and signs from the 18 case reports found in our literature search. The time lag from initial symptoms or thoracic insult to diagnosis ranged from a few minutes to over 16 hours. Awake patients (unlike ventilated patients) manifest the compensatory mechanisms,³ which have been clearly shown in animal studies (box 2). This leads to the disease being progressive and primarily respiratory (at least in appearance to the clinician) during the compensatory phase,³⁰ with hypoxemia being predominant.³ The evidence suggests that patients have progressive respiratory deterioration with final respiratory arrest, although rare cases of cardiac collapse are reported for which the mechanism remains unclear. Cyanosis and neurological sequelae are partially treatable with oxygen delivery to the patient and are hence variably present. The key to interpreting the early signs of hypoxia and respiratory distress is the degree of severity, but more importantly a pattern of relentless progression in a patient at risk of tension pneumothorax.^{3 51}

Ventilated patients

One case series of 71 patients with advanced TPT because of significant time delays before diagnosis (table 1) and 25 other case reports (box 4) were found in the literature search.

Diagnosis of TPT in ventilated patients rests with index of suspicion and recognition of supportive diagnostic features that the evidence shows to have more consistency than those found in awake patients. The development of tension is dependent on a pressure gradient between IPP and alveolar pressure. Ventilation will increase gas flow through the pleural defect, allowing more air to pass per unit time. This gives a more rapid IPP rise with earlier mechanical compressive effects and rapid progress to cardiorespiratory collapse, which will be further hastened by trauma and greater lung damage.^{3 78 79} Lung disease (ARDS/pulmonary haemorrhage/contusion) may also hasten decompensation by reducing lung compliance hence limiting lung collapse and thereby resulting in a greater IPP increase for any given gas volume.⁸⁰

A sudden fall in SpO₂ followed by hypotension (over a few minutes) was noted in Steier's large case series, is consistently found in the individual case reports and is also seen in reports from prehospital trauma care.^{36 81} There was only one report of slow development of TPT in a ventilated

Box 3 Symptoms and signs of tension pneumothorax from case reports in awake patients^{3 11 12 20 51-58}

Universal findings

- Chest pain
- Respiratory distress

Common findings (50%–75% cases)

- Tachycardia
- Ipsilateral decreased air entry

Inconsistent findings (<25% of cases)

- Low SpO₂
- Tracheal deviation
- Hypotension

Rare findings (about 10% cases)

- Cyanosis
- Hyper-resonance
- Decreasing level of consciousness
- Ipsilateral chest
 - Hyper-expansion
 - Hypo-mobility
- Acute epigastric pain
- Cardiac apical displacement
- Sternal resonance

patient—and this patient was partially self ventilating while un sedated.⁷⁵

Because of this sudden presentation, TPT in ventilated patients is more likely if the original pneumothorax is missed.³⁹ In the prehospital environment this means a low threshold for performing tube or finger thoracostomy must be maintained in ventilated trauma patients.^{36 81 82} In hospital, prevention depends on early diagnosis of pneumothorax. Although the chest radiological signs signs may be subtle (box 5) other alternatives for diagnosis exist such as: computed tomography,⁸³ or ultrasound⁸⁴ (with a negative predictive value as high as 100%).⁸⁵

The risk factors for pneumothorax misdiagnosis in ICU include: ventilation, unusual radiological location of pneumothorax, changed patient mental status, and lack of senior medical cover at the time of original presentation.⁸⁷⁻⁸⁹

Table 1 Signs of advanced tension pneumothorax (TPT) in ventilated patients—case series of 71 patients.²⁷

Signs of advanced TPT	Percentage
Subcutaneous emphysema	100
Tachycardia	95
Decreased breath sounds	87
Hyper-resonance	85
Systolic BP<90	81
Cyanosis	75
Low paO ₂	70
Tracheal deviation	60

Box 4 Signs of tension pneumothorax in ventilated patients—individual case reports^{1 10 28 49 50 59–77}

Universal findings

- Rapid onset
- Immediate and progressive decrease in arterial and mixed venous SpO₂
- Immediate reduction in cardiac output +/- BP

Common findings (each in about 33% of cases)

- High ventilation pressures
- Ipsilateral chest:
 - Hyper-expansion
 - Hypomobility
 - Decreased air entry

Inconsistent findings (each in about 20% of cases)

- Surgical emphysema
- Venous distension

Rarities that might make diagnosis difficult include bilateral TPT and loculated TPT causing cardiorespiratory compromise. The latter may occur if lung is bound to the chest wall by adhesions, (from parenchymal or pleural disease) so that it appears partially expanded on chest radiography.^{21 24 39 56 76 80}

MEDIASTINAL DEVIATION

Review articles of the chest radiological findings in TPT,^{24 90} and case reports with chest radiographs^{3 10 11 52–54 58} note mediastinal shift to be an inconsistent finding, which can also occur in the absence of TPT.^{10 24 52 75 91 92} The most severe mediastinal displacement described was in a 5 year old boy,⁵³ which is in keeping with Rutherford's³⁰ comments (and chest radiographs showing) the mobility of the mediastinum in children and immature monkeys in contrast with the immobility in adults and goats. Children may therefore suffer more rapid TPT development because of the more mobile mediastinum.³⁰

Even with chest radiological evidence of mediastinal displacement the trachea is often noted to be central—or unmentioned upon in the case report.^{3 10 11 52–54 58 93} Tracheal displacement is not commented upon in any of the animal experiments. Tracheal displacement was absent in all of the patients in a series of 108 suspected TPTs treated by paramedics with needle decompression³⁸ and present in only 1% of those receiving needle decompression by flight nurses in another series.⁹⁴ Even when tracheal displacement is known to be present the odds ratio for experienced physicians to correctly diagnose this has been shown to be no better than 50:50—that is, the same as tossing a coin.¹⁴ We suggest that tracheal displacement is neither diagnostic of TPT nor does its absence exclude the diagnosis.

CHEST SIGNS

Changing signs of chest mobility and expansion are poorly described features of this disease only being mentioned in one text reviewed⁷ of five major titles.^{5–9}

Ipsilateral hyper-expansion has been noted rarely in awake case reports although slightly more often in ventilated case reports but is to be expected when considering the pathophysiology. It occurs as a result of the maximal

Box 5 Chest radiological signs of pneumothorax

Ipsilateral

- sharp lung edge running parallel to the chest wall
- lucency
- deep lateral costo-phrenic angle (deep sulcus sign)⁸⁶
- abdominal quadrant hyperlucency

inspiratory efforts required to overcome the developing positive IPP with a transiently negative IPP for inspiration in awake patients. It also occurs as a result of the steadily expanding thoracic volume in both awake and ventilated patients. Ipsilateral chest hyper-expansion is consistently present on chest radiographs taken of TPT (evidenced by individual case reports,^{3 52 53 70 75} reviews of chest radiography in TPT,^{24 90} and animal experiments³⁰) and was well shown in the awake animal experiments.

Ipsilateral chest hypo-mobility may occur as a result of pain—pleuritic or from associated rib fractures. It will also occur in the latter stages as a result of movement limitation secondary to the hyper-expansion.

Contralateral hyper-mobility is expected in awake cases as part of the compensation to generate highly negative IPPs—and this is well shown in the awake animal studies.

It may be that the infrequent description of chest expansion and mobility signs may be attributable to the subtlety in their identification. Elicitation of these signs requires careful, longitudinal examination of the chest from the height of the patient, which is difficult in a critical clinical situation that potentially requires rapid management.

Other potential chest signs include ipsilateral added sounds, such as wheeze or crackles,³ the well described findings of decreased air entry with or without increased percussion note, sternal resonance, and displaced apex beat.

CARDIAC SIGNS

In awake patients only two case reports had early hypotension, which may have been related to TPT but one of these had significant other trauma as a possible explanation. Four patients developed hypotension between three and 16 hours after their initial insult, which had other potential causative factors (haemothorax, localised ventral TPT, myocardial ischaemia, bradycardia) in all but one. BP was preserved in all other 12 cases with tachycardia being noted in eight and no heart rate data in 10.

There was no evidence of great vessel obstruction in awake patients,³⁰ who (through respiratory decompensation), may lack the capacity to achieve this level of IPP. Even when mediastinal shift is evident on chest radiography there is evidence of poor correlation with haemodynamic status.⁹³ Decreased BP or cardiac output are not common in awake patients and if found are likely to be attributable to other pathology or the hypoxic effects of TPT when it is in the advanced stages. TPT induced hypotension and/or rapidly decreasing SpO₂ is almost certainly pre-terminal.

Distension of neck veins is infrequently commented upon in both case reports and animal experiments, will only occur if there is no coexisting hypovolaemia and even then may be obscured by a cervical collar in trauma.

The decreased cardiac output in ventilated patients is consistent, early, and progressive. It is probably attributable to a combination of hypoxaemia,^{3 95} diminished blood flow through the collapsed lung,^{30 96 97} reduced venous return, and possibly great vessel/ventricular compression.^{47 49} Hypotension is pre-terminal^{3 30 49} and if TPT is not recognised and

treated urgently it is especially likely in ventilated patients to lead to sudden cardiac arrest.^{1 10}

CHEST RADIOGRAPHY IN TENSION PNEUMOTHORAX

Box 6 lists the chest radiological findings of tension pneumothorax. It should be noted that merely looking for mediastinal shift may not give conclusive differentiation of a TPT from a simple pneumothorax as this may occur in both conditions.^{24 34} In rare cases a loculated anterior TPT may exist that will only be visible on a lateral chest radiography or computed tomography.^{56 76 98 99}

The use of chest radiography to diagnose TPT has been associated with a fourfold increase in mortality based on the evidence from two studies in ventilated patients who waited between 30 minutes and eight hours for the chest radiograph.^{27 87} This high mortality was almost certainly attributable to the prolonged delay when the diagnosis was not made clinically or the chest radiograph was misinterpreted. The high conversion rate from pneumothorax to TPT in ventilated patients²⁷ further alerted clinicians to diagnose the condition earlier and treat before chest radiograph. The concept of; “the chest radiograph that should never have been taken,” has since been emphasised.^{3 5 35 100}

However, other evidence does not support this blanket statement. Chest radiography has been used (apparently without adverse outcome) to confirm TPT diagnosis in stable patients with equivocal diagnostic signs,^{52 53 70 75} so avoiding unnecessary morbidity and potentially offering alternative diagnoses.¹⁰¹ Chest radiography has also diagnosed unexpected TPT in unstable ventilated patients when a lack of other signs has prevented earlier diagnosis.^{63 65 66 71 77} It has been suggested that in awake patients in the emergency department a chest radiograph is obtained before tube thoracostomy thus sparing some patients the morbidity and discomfort of a potentially unnecessary procedure.^{18 102} In a retrospective analysis of 176 emergency department presentations of pneumothorax none of the 30 patients identified with mediastinal shift on chest radiography, had decompressed while awaiting tube thoracostomy, and two further patients in this series with haemodynamic compromise were decompressed before chest radiography.⁹³

Ultimately whether or not radiological confirmation of the diagnosis takes place must be a balanced risk of the time taken to obtain the test, and the expected clinical course. We suggest that in ventilated patients decompression should still precede chest radiography where the diagnosis of TPT is suspected. However, in awake patients we suggest that a chest radiograph precedes decompression unless the patient is unstable and potentially in the advanced stages of tension—or an immediate chest radiograph is unavailable. If a chest radiograph is not immediately available box 7 lists our recommendations for immediate decompression, (which

Box 6 Chest radiological findings in tension pneumothorax^{24 80 90}

Ipsilateral hyper-expansion

- hemi-diaphragmatic depression
- increased rib separation
- increased thoracic volume

Mediastinal pressure

- ipsilateral flattening of heart border
- contralateral mediastinal deviation

Box 7 Recommendations for immediate chest decompression in awake patients with tension pneumothorax as the suspected cause

Chest radiograph not immediately available and:

- SpO₂<92% on O₂
- Systolic BP<90 mm Hg
- Respiratory rate<10
- Decreased level of consciousness on O₂
- Cardiac arrest
 - bilateral finger or tube thoracostomy
 - not needle thoracocentesis

are in partial agreement with previous indications for prehospital needle decompression).³⁸ When the chest radiograph is taken a practitioner capable of urgent decompression should stay with the patient.

NEEDLE DECOMPRESSION

Emergency needle decompression followed by tube thoracostomy is widely advocated^{5 103} but it must be appreciated that it has an associated morbidity (box 8).

It may also be ineffective with decompression only being adequately achieved on tube thoracostomy.¹⁵⁻²² Barton⁹⁴ found that of 106 patients treated with tube thoracostomy by prehospital flight nurses 38% had been attributable to failure of clinical improvement with needle decompression. Box 9 lists the potential causes of needle decompression failure.

No studies exist showing the sensitivity of needle decompression²¹ and yet it is regularly taught as a “rule out” investigation. This assumption of needle thoracocentesis having a 100% negative predictive value is clearly dangerous in the treatment of severely compromised patients, especially those with a potentially reversible cause of cardiac arrest.¹

Despite the problems with needle decompression, it is a technique used commonly in the prehospital environment,^{32 37 38 82} has proved safe and therapeutic,^{18 36 37 94} and has been shown to lead to shorter on-scene times compared with tube thoracostomy.⁹⁴ However, some feel that the number of needle decompressions far exceeds the incidence of TPT and should be limited—especially in the prehospital environment.^{18 32 102} An examination of needle decompression in 19 trauma patients, admitted to an American level 1 trauma centre found that only four patients had evidence of

Box 8 Morbidity associated with needle decompression^{18 94 102 104 105}

- Unnecessary pain/discomfort for the patient
 - if procedure was not required
- Pneumothorax
 - with potential to tension later—especially if ventilated
- Cardiac tamponade
- Life threatening haemorrhage
- Loculated intrapleural haematoma
- Atelectasis
- Pneumonia

Box 9 Potential causes of needle decompression failure^{17 18 20 21 94}

- Obstruction by:
 - blood
 - tissue
 - kinking
- Missing a localised tension pneumothorax
- Inability to drain a large air leak
- Requirement for repeated needle decompression

pleural air leak and only two of these had tension physiology with just one being adequately decompressed by needle.¹⁸

Needle decompression in the 2nd/3rd intercostal space (ICS), mid-clavicular line (MCL) is easy to access, but entails penetration of pectoral muscles and a variable quantity of subcutaneous tissue with or without oedema and subcutaneous emphysema. A standard 14 gauge (4.5 cm) cannula may not be long enough to penetrate parietal pleura,^{15 16 19 22} with up to one third of trauma patients having a chest wall thickness greater than 5 cm in the 2nd ICS MCL.¹⁰⁶ This will lead to treatment failure and diagnostic confusion, although using the trocar instead (7 cm) may negate this problem and prevent kinking.²² The use of the 4th or 5th ICS in the mid-axillary line may be safer¹⁰⁵ and has been recommended by ATLS⁵ as it contains less fat and avoids large muscles. Unfortunately this site may have an increased risk of lung damage in the supine patient, as gas collects at the highest point and adhesions are most likely in more dependent parts of the lung.¹⁰⁷

A syringe filled with sterile saline attached to the cannula may help confirm pleural penetration^{91 108} and an Asherman chest seal may help stabilise the cannula preventing displacement or kinking.¹⁰⁹ Some authors advocate the attachment of a flutter valve to the cannula¹⁰³ but if attached wrongly this may cause re-tension.¹¹⁰

TUBE THORACOSTOMY

The rapid potential deterioration of TPT means that some suggest blind tube thoracostomy upon disease suspicion or after needle decompression.^{5 6} This is arguably the treatment of choice, ensuring maximal pleural cavity evacuation and lung re-expansion.⁹⁴ It has been shown to be safe and effective in trained physician led prehospital trauma care.^{36 82 94} In Coats' series³⁶ 57 patients with TPT who had received physician led blind needle decompression or tube thoracostomy in prehospital trauma care showed an improvement in BP and SpO₂, with one case of infection and one episode of tube misplacement, which was recognised and corrected on scene. Schmidt⁸² noted no cases of re-tension after tube thoracostomy in his series of 76 patients. This approach inevitably has an over-diagnosis rate with 15 of 76 patients in whom tube thoracostomy was performed, having no haemopneumothorax or pneumothorax on arrival at the trauma centre.⁸²

The reported complication rate of tube thoracostomy varies from 4% to 30% with only 1% occurring from insertion.^{111–114} Complications (box 10) are probably less common now that use of the trocar has been abandoned in place of blunt dissection,¹¹⁵ although fatalities may still occur.¹¹⁶

Many tube thoracostomy complications are related to the tube itself—re-tension, haematoma, infection, subcutaneous placement.⁹⁴ Simple thoracostomy without tube placement is an effective alternative, (in ventilated patients) that avoids

Box 10 Complications of tube thoracostomy^{117 118}

- Death
- Haemorrhage (most commonly intercostal artery damage)
- Bronchopleural fistula
- Subcutaneous tube placement
- Intra-peritoneal tube placement
- Infection
- Damage to
 - Lung parenchyma
 - Mediastinal contents
 - Neurovascular bundles
- Myocardium

these complications. It also decreases time to decompression, decreases on scene time in prehospital care, and enables repeat 360° finger sweep in the event of patient deterioration.⁸¹ The latter will permit palpation of the lung to check for continued re-expansion and confirm the patency of pleural communication. The procedure simply entails a thoracostomy in the 4th or 5th ICS mid-axillary line, ensuring good finger access to the pleural cavity to permit decompression.

SUMMARY

Any pleural injury communicating with the atmosphere via a one way valve that opens on inspiration and closes on expiration will lead to an expanding pneumothorax. Defining the point of tension without IPP measurements is more difficult and it might be argued that “expanding pneumothorax” is a more appropriate term. For clinical purposes however we would suggest that a pneumothorax is considered to be under tension when it results in “significant respiratory or haemodynamic compromise (the latter especially in ventilated patients) that reverses on decompression alone”—that is, without chest drain placement. A continual egress of air throughout the respiratory cycle will add to the diagnostic certainty—unlike a brief hiss during one phase of the respiratory cycle.

The true incidence of TPT is unknown but it is more common in ventilated than awake patients and possibly most common in ventilated patients with visceral pleural injury from chest trauma.⁵

The differences in the pathophysiology and presentation of TPT between awake and ventilated patients have been mentioned by other authors^{3 50} but deserve increased emphasis. TPT should not be taught as if it was a single entity. The natural history of the disease is of progressive hypoxia with a variety of cardiorespiratory compensatory mechanisms, which are well seen in awake patients but are blunted by ventilation and sedation.

In ventilated patients TPT presents rapidly with consistent signs of respiratory and cardiac compromise. In contrast awake patients show a greater variability of presentations, which are generally more progressive, with slower decompression. In both groups the general signs are more consistent than the lateralising signs and we suggest that the clinician attempts first to diagnose the condition and then to lateralise it. In trauma if the diagnosis of TPT is in doubt these lateralising signs may also help to differentiate TPT from other causes of severe respiratory embarrassment such as flail chest and pulmonary contusion. We feel that tracheal deviation and neck vein distension should be de-emphasised, if not abandoned altogether, as taught signs

of the disease. The clinician should also be aware of the pre-terminal signs in either group of patients and the fact that a ventilated patient is more likely to suffer a cardiac arrest in contrast with the awake patient who is more likely to suffer a respiratory arrest. We suggest the signs and symptoms listed in boxes 11 and 12 are adapted as the way of diagnosing TPT in both of these groups.

Once diagnosed treatment should begin with immediate high flow oxygen to maximise oxygenation and provide an oxygen reservoir for further resuscitation. Upright positioning may improve the patients condition and be beneficial for survival,^{11 30} but should not be done in multi-trauma or other cases of suspected spinal injury.

Needle decompression still has a place and is potentially a life saver, but its indiscriminate use should be discouraged. Its potential for failure, the reasons for this, and alternative approaches should be emphasised in teaching. In awake patients it should be performed when there is specific evidence of decompensation (box 6) although immediate tube thoracostomy is preferred. Otherwise we would encourage that the clinician obtains a chest radiograph to confirm the diagnosis and lateralise the disease while being prepared to perform needle decompression should the patient decompensate.

In ventilated ICU patients needle decompression has a place in the immediate treatment of suspected TPT but where

Box 11 Diagnosis of tension pneumothorax in ventilated patients

Rapid disease progression

Early reliable

- Decrease in SpO₂—immediate
- Decrease in Q
- Decrease in BP
- Tachycardia

Others

- Increased ventilation pressure
- Surgical emphysema

Disease lateralisation

- Ipsilateral
 - Hyper-resonance
 - Decreased breath sounds
 - Chest hyper-expansion
 - Chest hypo-mobility
 - Added sounds

Inconsistent

- Cyanosis
- Distended neck veins
- Tracheal deviation

Monitor dependent^{32 50 95}

- Increasing CVP
- Increasing pulmonary arterial pressure
- ECG data
- Decreased mixed venous oxygen saturations

Box 12 Diagnosis of tension pneumothorax in awake patients

Reliable and early

- Pleuritic chest pain
- Air hunger
- Respiratory distress
- Tachypnoea
- Tachycardia
- Falling SpO₂
- Agitation

Disease lateralisation—ipsilateral

- Hyper-expansion
- Hypo-mobility
- Hyper-resonance
- Decreased breath sounds
- Added sounds—crackles/wheeze

Disease lateralisation—contralateral

- Hyper-mobility

Pre-terminal^{3 49 54}

- Decreasing respiratory rate
- Hypotension
- Decreasing SpO₂
- Decreasing level consciousness

Inconsistent

- Tracheal deviation
- Distended neck veins

possible immediate tube thoracostomy should be encouraged as the treatment of choice. In a ventilated patient with significant chest trauma and undiagnosed shock or cardiac arrest (for example, in the prehospital or immediate resuscitation room phase) we would advocate formal thoracostomy (finger or tube) as an immediate treatment. However, it should also be noted that in unskilled hands the morbidity of thoracostomy is not insignificant (box 10).

Tension pneumothorax is a potential killer and the possibility of it should always be borne in mind. Its diagnosis however may be difficult even in the hands of experienced clinicians and the decisions regarding the best treatment option and its timing require critical decisions to be made. It is hoped that this review will help with that process and we encourage the submission and publication of further detailed case reports to build up the knowledge base—especially in the more variable disease that occurs in awake patients.

ACKNOWLEDGEMENTS

The authors would like to thank Mr Tim Coats, Professor of Emergency Medicine (Barts and the London, Queen Mary's School of Medicine, University of London) who provided invaluable support and advice during preparation of the final manuscript.

CONTRIBUTORS

Mr Leigh-Smith initiated the review, performed the original literature search, prepared the first draft, and collated the two authors drafts. Dr Harris performed a separate literature search and produced an independent draft. Both authors contributed to the final draft. Mr Leigh-Smith is the guarantor of this paper.

Authors' affiliations

S Leigh-Smith, Defence Medical Services, UK
T Harris, Intensive Care Unit, Alfred Hospital, Melbourne, Australia

Funding: none.

Conflicts of interest: none declared.

REFERENCES

- Rojas R, Wasserberger J, Balasubramaniam S. Unsuspected tension pneumothorax as a hidden cause of unsuccessful resuscitation. *Ann Emerg Med* 1983;**12**:411–12.
- Fischer H, Masel H. Spontaneous pneumothorax and tension pneumothorax as causes of sudden death (author's translation). *Z Rechtsmedizin—Journal of Legal Medicine* 1978;**81**:223–6.
- Barton, ed. Tension pneumothorax. *Curr Opin Pulm Med* 1999;**5**:269–74.
- ALSG. *Advanced life support manual*. 4th ed. London: Resuscitation Council (UK), 2001.
- ATLS. *Advanced trauma life support*. 6th ed. Chicago: American College of Surgeons, 1997.
- Tintinalli J, Kelen G, Stapczynski J. *Emergency medicine—a comprehensive study guide*. 5th ed. New York: McGraw Hill, 2000.
- Greaves I, Porter K, Ryan J. *Trauma care*. London: Arnold, 2001.
- Rosen P, Barkin R, Danzl D, et al. *Emergency medicine—concepts and clinical practice*. 4th ed. St Louis: Mosby, 1999.
- Wyatt J, Illingworth R, Clancy M, et al. *Oxford handbook of accident and emergency medicine*. Oxford: Oxford University Press, 1999.
- Watts BL, Howell MA. Tension pneumothorax: a difficult diagnosis. *Emerg Med J* 2001;**18**:319–20.
- Holloway VJ, Harris JK. Spontaneous pneumothorax: is it under tension? *J Accid Emerg Med* 2000;**17**:222–3.
- Hollins GW, Beattie T, Harper I, et al. Tension pneumothorax: report of two cases presenting with acute abdominal symptoms. *J Accid Emerg Med* 1994;**11**:43–4.
- Dunlop MG, Beattie TF, P. G. P. Clinical assessment and radiography following blunt chest trauma. *Arch Emerg Med* 1989;**6**:125–7.
- Spiteri MA, Cook DG, Clark SW. Reliability of eliciting physical signs in examination of the chest. *Lancet* 1988;**i**:873–5.
- Britten S, Palmer SH. Chest wall thickness may limit adequate drainage of tension pneumothorax by needle thoracocentesis. *J Accid Emerg Med* 1996;**13**:426–7.
- Britten S, Palmer SH, Snow TM. Needle thoracocentesis in tension pneumothorax: insufficient cannula length and potential failure. *Injury* 1996;**27**:321–2.
- Conces DJ Jr, Tarver RD, Gray WC, et al. Treatment of pneumothoraces utilizing small caliber chest tubes. *Chest* 1988;**94**:55–7.
- Cullinane DC, Morris JA Jr, Bass JG, et al. Needle thoracostomy may not be indicated in the trauma patient. *Injury* 2001;**32**:749–52.
- Jenkins C, Sudheer PS. Needle thoracocentesis fails to diagnose a large pneumothorax. *Anaesthesia* 2000;**55**:925–6.
- Jones R, Hollingsworth J. Tension pneumothoraces not responding to needle thoracocentesis. *Emerg Med J* 2002;**19**:176–7.
- Mines D, Abbuhl S. Needle thoracostomy fails to detect a fatal tension pneumothorax [erratum appears in *Ann Emerg Med* 1993;**22**:1364]. *Ann Emerg Med* 1993;**22**:863–6.
- Pattison GT. Needle thoracocentesis in tension pneumothorax: insufficient cannula length and potential failure. *Injury* 1996;**27**:758.
- Driscoll P, Gwinutt C, Graham T. Chest and cardiac trauma. In: Skinner D, Swain A, Peyton R, eds. *Cambridge textbook of accident and emergency medicine*. Cambridge: Cambridge University Press, 1997:538.
- Teplick SK, Clark RE. Various faces of tension pneumothorax. *Postgrad Med* 1974;**56**:87–92.
- Saltet JF. Pneumothorax. *Lancet* 1979;**i**:671.
- Ludwig J, Kienzle GD. Pneumothorax in a large autopsy population. A study of 77 cases. *Am J Clin Pathol* 1978;**70**:24–6.
- Steier M, Ching N, Roberts EB, et al. Pneumothorax complicating continuous ventilatory support. *J Thorac Cardiovasc Surg* 1974;**67**:17–23.
- Yu PY, Lee LW. Pulmonary artery pressures with tension pneumothorax. *Can J Anaesth* 1990;**37**:584–6.
- Chen KY, Jerng JS, Liao WY, et al. Pneumothorax in the ICU: patient outcomes and prognostic factors. *Chest* 2002;**122**:678–83.
- Rutherford RB, Hurt HH Jr, Brickman RD, et al. The pathophysiology of progressive, tension pneumothorax. *J Trauma* 1968;**8**:212–27.
- Seaton D. Pneumothorax. In: Seaton A, Leitch A, eds. *Crofton and Douglas's respiratory diseases*. 5th ed. Oxford: Blackwell Science, 2000:1182–211.
- Bjerke H. Tension pneumothorax. *Emergicine Specialities*, 2002. (<http://www.emedicine.com/med/topic2793.htm>).
- Gustman P, Yerger L, Wanner A. Immediate cardiovascular effects of tension pneumothorax. *Am Rev Respir Dis* 1983;**127**:171–4.
- Fraser R, Pare J. Tension pneumothorax and hydrothorax. In: *Diagnosis of diseases of the chest*. Philadelphia: W B Saunders, 1978:598.
- Light RW. Tension pneumothorax. *Intensive Care Med* 1994;**20**:468–9.
- Coats TJ, Wilson AW, Xeropotamos N. Pre-hospital management of patients with severe thoracic injury. *Injury* 1995;**2**:581–5.
- Cameron PA, Fleit K, Kacan E, et al. Helicopter retrieval of primary trauma patients by a paramedic helicopter service. *Aust N Z J Surg* 1993;**63**:790–7.
- Eckstein M, Suyehara D. Needle thoracostomy in the prehospital setting. *Prehosp Emerg Care* 1998;**2**:132–5.
- Tocino IM, Miller MH, Fairfax WR. Distribution of pneumothorax in the supine and semirecumbent critically ill adult. *AJR Am J Roentgenol* 1985;**144**:901–5.
- Simmons D, Hemingway A, Richiuti N. Acute circulatory effects of pneumothorax in dogs. *J Appl Physiol* 1958;**12**:255–61.
- Hilton R. Some effects of artificial pneumothorax on the circulation. *J Pathol Bacteriol* 1925;**37**:1–8.
- Kilburn K. Cardiorespiratory effects of large pneumothorax in conscious and anaesthetised dogs. *J Appl Physiol* 1963;**18**:279–83.
- Maloney JV Jr, Schmutzer K, Raschke E. Paradoxical respiration and "pendelluft." *J Thorac Cardiovasc Surg* 1961;**41**:291–8.
- Stewart H, Bailey Jr R. The effect of unilateral spontaneous pneumothorax on the circulation in man. *J Clin Invest* 1940;**19**:321–6.
- Bennett RA, Orton EC, Tucker A, et al. Cardiopulmonary changes in conscious dogs with induced progressive pneumothorax. *Am J Vet Res* 1989;**50**:280–4.
- Carvalho P, Hilderbrandt J, Charan NB. Changes in bronchial and pulmonary arterial blood flow with progressive tension pneumothorax. *J Appl Physiol* 1996;**81**:1664–9.
- Hurewitz AN, Sidhu U, Bergofsky EH, et al. Cardiovascular and respiratory consequences of tension pneumothorax. *Bull Eur Physiopathol Respir* 1986;**22**:545–9.
- West J. *The mechanics of breathing*. *Respiratory physiology—the essentials*. 5th ed. Baltimore: Williams and Wilkins, 1995:31–50, 89–116.
- Barton ED, Rhee P, Hutton KC, et al. The pathophysiology of tension pneumothorax in ventilated swine. *J Emerg Med* 1997;**15**:147–53.
- Beards SC, Lipman J. Decreased cardiac index as an indicator of tension pneumothorax in the ventilated patient. *Anaesthesia* 1994;**49**:137–41.
- Leigh-Smith S, Davies G. Tension pneumothorax—eyes may be more diagnostic than ears. *Emerg Med J* 2003;**20**:1–2.
- Askins DC. Spontaneous tension pneumothorax during sexual intercourse. *Ann Emerg Med* 1984;**13**:303–6.
- Boon D, Llewellyn T, Rushton P. A strange case of a tension pneumothorax. *Emerg Med J* 2002;**19**:470–1.
- Friend KD. Prehospital recognition of tension pneumothorax. *Prehosp Emerg Care* 2000;**4**:75–7.
- Slay RD, Slay LE, Luehrs JG. Transient ST elevation associated with tension pneumothorax. *JACEP* 1979;**8**:16–18.
- Vermeulen EG, Teng HT, Boxma H. Ventral tension pneumothorax. *J Trauma* 1997;**43**:975–6.
- Wilkinson D, Moore E, Wither P, et al. ATLS on the ski slopes—a steamboat experience. *J Trauma* 1992;**32**:448–51.
- Werne CS, Sands MJ. Left tension pneumothorax masquerading as anterior myocardial infarction. *Ann Emerg Med* 1985;**14**:164–6.
- Baldwin LN. Mixed venous oximetry in the diagnosis of tension pneumothorax. *Anaesthesia* 1995;**50**:181–2.
- Cohen Y, Laker M. Tension pneumothorax from inappropriate oxygen administration. *Harefuah* 1993;**124**:549–51, 599.
- Gambrell VL. Diagnosis and treatment of tension pneumothorax under anesthesia: a case report. *AANA Journal* 2002;**70**:21–4.
- Hardy MJ, Huard C, Lundblad TC. Bilateral tension pneumothorax during jet ventilation: a case report. *AANA Journal* 2000;**68**:241–4.
- Janssens U, Koch KC, Graf J, et al. Severe transmural ischemia in a patient with tension pneumothorax. *Crit Care Med* 2000;**28**:1638–41.
- Monterrubio Villar J, Fernandez Berges D, Alzugaray Fraga RJ, et al. [ST elevation and tension pneumothorax]. *Rev Esp Cardiol* 2000;**53**:467–70.
- Strizik B, Forman R. New ECG changes associated with a tension pneumothorax: a case report. *Chest* 1999;**115**:1742–4.
- Wells L, Aves T. Pneumothorax presenting as ischaemia of the hand. *Anaesthesiology* 1995;**82**:586–7.
- Woodcock TE, Murray S, Ledingham IM. Mixed venous oxygen saturation changes during tension pneumothorax and its treatment. *Anaesthesia* 1984;**39**:1004–6.
- Plewa MC, Ledrick D, Sferra JJ. Delayed tension pneumothorax complicating central venous catheterization and positive pressure ventilation. *Am J Emerg Med* 1995;**13**:532–5.
- Cridde LM. An unusual case of tension pneumothorax. *Am J Crit Care* 1995;**4**:314–18.
- Hosteier MA, Davis CO. Bilateral localized tension pneumothoraces refractory to needle decompression. *Pediatr Emerg Care* 1999;**15**:322–4.
- Engoren M, de St Victor P. Tension pneumothorax and contralateral presumed pneumothorax from endobronchial intubation via cricothyroidotomy. *Chest* 2000;**118**:1833–5.
- Rai A, Iftikhar S. Tension pneumothorax complicating diagnostic upper endoscopy: a case report. *Am J Gastroenterol* 1999;**94**:845–7.
- Seher G, Janda A. Tension pneumothorax as anaesthesia complication—report on 3 cases (author's translation). *Anaesthesist* 1975;**24**:183–4.
- Laisley RS, Aps C. Tension pneumothorax and pulse oximetry. *Br J Anaesth* 1991;**66**:250–2.
- Connolly JP. Hemodynamic measurements during a tension pneumothorax. *Crit Care Med* 1993;**21**:294–6.
- Ross IB, Fleiszer DM, Brown RA. Localized tension pneumothorax in patients with adult respiratory distress syndrome. *Can J Surg* 1994;**37**:415–19.
- Loer S, Fritz KW. [Life threatening tension pneumothorax after puncture of the subclavian vein and dislocation of thoracic drainage]. *Anaesthesiol Reanim* 1994;**19**:137–8.
- Mariani PJ, Sharma S. Iatrogenic tension pneumothorax complicating outpatient Heimlich valve chest drainage. *J Emerg Med* 1994;**12**:477–9.
- Tan EC, van der Vliet JA. [Delayed (tension) pneumothorax after placement of a central venous catheter]. *Ned Tijdschr Geneesk* 1999;**143**:1872–5.

- 80 **Gobien RP**, Reines HD, Schabel SI. Localized tension pneumothorax: unrecognized form of barotrauma in adult respiratory distress syndrome. *Radiology* 1982;**142**:15-19.
- 81 **Deakin CD**, Davies G, Wilson A. Simple thoracostomy avoids chest drain insertion in prehospital trauma. *J Trauma* 1995;**39**:373-4.
- 82 **Schmidt U**, Stalp M, Gerich T, *et al*. Chest tube decompression of blunt chest injuries by physicians in the field: effectiveness and complications. *J Trauma* 1998;**44**:98-101.
- 83 **Bourgouin P**, Cousineau G, Lemire P, *et al*. Computer tomography used to exclude pneumothorax in bullous lung disease. *J Can Assoc Radiol* 1985;**36**:341-2.
- 84 **Chan SS**. Emergency bedside ultrasound to detect pneumothorax. *Acad Emerg Med* 2003;**10**:91-4.
- 85 **Lichtenstein DA**, Menu Y. A bedside ultrasound sign ruling out pneumothorax in the critically ill. Lung sliding. *Chest* 1995;**108**:1345-8.
- 86 **Gordon R**. The deep sulcus sign. *Radiology* 1980;**136**:25-7.
- 87 **Kollef MH**. Risk factors for the misdiagnosis of pneumothorax in the intensive care unit. *Crit Care Med* 1991;**19**:906-10.
- 88 **Kollef MH**. Hemodynamic measurements during a tension pneumothorax. *Crit Care Med* 1994;**22**:896.
- 89 **Kollef MH**. The effect of an increased index of suspicion on the diagnosis of pneumothorax in the critically ill. *Mil Med* 1992;**157**:591-3.
- 90 **Greene R**, McCloud TC, Stark P. Pneumothorax. *Semin Roentgenol* 1977;**12**:313-25.
- 91 **Light R**. *Textbook of respiratory medicine*. Philadelphia: WB Saunders, 1988.
- 92 **Kupferschmid JP**, Carr T, Fonger JD, *et al*. Chronic tension pneumothorax mimicking tension bullae. Use of video-assisted thoracoscopy for diagnosis. *Chest* 1993;**104**:1913-14.
- 93 **Clark S**. Is mediastinal shift always an emergency? ACEM winter symposium, Melbourne, 2003.
- 94 **Barton ED**, Epperson M, Hoyt DB, *et al*. Prehospital needle aspiration and tube thoracostomy in trauma victims: a six-year experience with aeromedical crews. *J Emerg Med* 1995;**13**:155-63.
- 95 **Gilbert TB**. Hemodynamic consequences of tension pneumothorax. *Crit Care Med* 1993;**21**:1981-2.
- 96 **Richards DW Jr**, Riley C, Hiscock M. Cardiac output following artificial pneumothorax in man. *Arch Intern Med* 1932;**49**:994-1006.
- 97 **Weiss R**. Ueber die Durchblutung der Kollapslung beim experimentellen Pneumothorax. *Z Exp Med* 1926;**53**.
- 98 **Collins JA**, Samra GS. Failure of chest X-rays to diagnose pneumothoraces after blunt trauma. *Anaesthesia* 1998;**53**:74-8.
- 99 **Jantsch H**, Winkler M, Pichler W, *et al*. [Pneumothorax in intensive care patients. The value of tangential views]. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr* 1990;**153**:275-7.
- 100 **Baumann MH**, Sahn SA. Tension pneumothorax: diagnostic and therapeutic pitfalls. *Crit Care Med* 1993;**21**:177-9.
- 101 **Coren ME**, Rosenthal M, Bush A. Congenital diaphragmatic hernia misdiagnosed as tension pneumothorax. *Pediatr Pulmonol* 1997;**24**:119-21.
- 102 **Etoch SW**, Bar-Natan MF, Miller FB, *et al*. Tube thoracostomy. Factors related to complications. *Arch Surg* 1995;**130**:521-5, 525-6.
- 103 **Bowman J**. Pneumothorax, tension and traumatic. <http://www.emedicine.com/emerg/topic470.htm>, 2002.
- 104 **Butler KL**, Best IM, Weaver WL, *et al*. Pulmonary artery injury and cardiac tamponade after needle decompression of a suspected tension pneumothorax. *J Trauma* 2003;**54**:610-11.
- 105 **Rawlins R**, Brown KM, Carr CS, *et al*. Life threatening haemorrhage after anterior needle aspiration of pneumothoraces. A role for lateral needle aspiration in emergency decompression of spontaneous pneumothorax. *Emerg Med J* 2003;**20**:383-4.
- 106 **Marinero J**, Kenny C, Smith S, *et al*. Needle thoracostomy in trauma patients: what catheter length is adequate? *Acad Emerg Med* 2003;**10**:495.
- 107 **Goodman L**, Putman C. *Intensive care radiology/imaging of the critically ill*. 2nd ed. Philadelphia: W B Saunders, 1982:99-100.
- 108 **Light R**. *Pneumothorax/Pleural disease*. 3rd ed. Baltimore: Williams and Wilkins, 1995:242-77.
- 109 **Allison K**, Porter KM, Mason AM. Use of the Asherman chest seal as a stabilisation device for needle thoracostomy. *Emerg Med J* 2002;**19**:590-1.
- 110 **Mainini SE**, Johnson FE. Tension pneumothorax complicating small-caliber chest tube insertion. *Chest* 1990;**97**:759-60.
- 111 **Millikan J**, Moore E, Steiner E, *et al*. Complications of tube thoracostomy for acute trauma. *Am J Surg* 1980;**140**:738-41.
- 112 **Daly RC**, Mucha P, Pairloro PC, *et al*. The risk of percutaneous chest tube thoracostomy for blunt thoracic trauma. *Ann Emerg Med* 1985;**14**:865-70.
- 113 **Chan L**, Reilly KM, Henderson C, *et al*. Complication rates of tube thoracostomy. *Am J Emerg Med* 1997;**15**:368-70.
- 114 **Bailey RC**. Complications of tube thoracostomy in trauma. *J Accid Emerg Med* 2000;**17**:111-14.
- 115 **Hyde J**, Sykes T, Graham T. Reducing morbidity from chest drains. *BMJ* 1997;**314**:914-15.
- 116 **Muckart DJ**, Aitchison JM. Fatality from blind intubation of suspected tension pneumothorax. *Injury* 1989;**20**:175-6.
- 117 **Tang AT**, Velissaris TJ, Weeden DF. An evidence-based approach to drainage of the pleural cavity: evaluation of best practice. *J Eval Clin Pract* 2002;**8**:333-40.
- 118 **Phillips J**. The chest (pneumothorax, haemothorax, effusions and empyema). *The Internet Journal of Thoracic and Cardiovascular Surgery* 2003;**5**. (<http://www.ispub.com/ostia/index.php?xmlfilepath=journals/ijtcvs/vol5n2/chest.xml>).



Tension pneumothorax—time for a re-think?

S Leigh-Smith and T Harris

Emerg Med J 2005 22: 8-16

doi: 10.1136/emj.2003.010421

Updated information and services can be found at:

<http://emj.bmj.com/content/22/1/8.full.html>

These include:

References

This article cites 94 articles, 26 of which can be accessed free at:

<http://emj.bmj.com/content/22/1/8.full.html#ref-list-1>

Article cited in:

<http://emj.bmj.com/content/22/1/8.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://journals.bmj.com/cgi/ep>